

Endocrinopathies after Traumatic Brain Injury

Susan R. Rose, MD
Professor
Cincinnati Children's Hospital Medical Center
University of Cincinnati



6th Annual Northern Kentucky TBI Conference
March 23, 2012
www.bridgesnky.org

- Supported by
 - Pfizer, Inc
 - Investigator-Initiated Research grant
 - General Clinical Research Center, NIH
 - #M01 RR08084
 - Nat'l Center for Research Resources, NIH
 - USPHS Grant #UL1 RR026314

Traumatic Brain Injury (TBI) in children

- TBI is leading cause of death & disability in children.
- Incidence
 - Infants (inflicted injury) 25/ 100,000
 - Children 180/ 100,000
 - Peak age 15-24y, 250/ 100,000
- About 6 million Americans have lifelong disability after TBI

Causes of TBI vary with age

- Infants-- inflicted injuries
- Toddlers—falls
- School-aged children--pedestrian or sports-related injuries
- Adolescents--motor vehicle crashes

Costs of TBI

- Cost for care of one person can be \$2 million
- In 1995 in USA
 - costs & lost productivity from TBI= \$56 billion
- Consequences:
 - Neurologic
 - Cognitive
 - Behavior deficits
 - Social
 - Economic
 - Endocrine

However

- Endocrine outcomes of TBI are under-recognized
 - Not mentioned in
 - 1998 NIH Consensus Conference Statement
 - most TBI websites (ninds, niccyd, cdc)

Outline



- Mechanisms of injury
- Endocrine deficiencies
- Endocrine studies after TBI
- CCHMC study
- Consensus conference
- Conclusions



Types of CNS injury

- **Open Head Injury**
 - Penetration of the skull–focal
- **Closed Head Injury**
 - Focal & diffuse damage to axons
- **Deceleration Injuries**
 - Different parts of brain move at different speeds.
 - diffuse axonal shearing, contusion, brain swelling.
 - axons stretched until torn.
- **Hypoxia**
 - irreversible brain injury from anoxia (no oxygen)
- **Stroke**
 - cerebral vascular accident, cell death in area deprived of blood
 - bleeding in or over brain (hemorrhage or hematoma)

Coup and Contra-coup





Mechanism of injury

Ropper AH,
N Engl J Med 2007

Mechanical injury to hypothalamus & pituitary in TBI

- Anatomic mechanisms responsible for injury
 - direct damage to pituitary stalk
 - disruption of vascular supply
 - indirect effects on function due to poor perfusion from cerebral edema or shock
 - pituitary antibodies after injury




Location of injury

- If hypothalamus
 - low anterior pituitary hormones
 - normal response to releasing hormones.
- If lower pituitary stalk or anterior lobe
 - low anterior pituitary hormones
 - no response to releasing hormones
 - High prolactin
 - SIADH or DI
- Most severe injuries
 - damage both hypothalamus & pituitary
 - mixed endocrine picture.

Yuan 1991

Outline

- Mechanisms of injury
- Endocrine deficiencies
- Endocrine studies after TBI
- CCHMC study
- Consensus conference
- Conclusions



Common symptoms of Hypopituitarism

- Not specific for any one deficiency
 - Weakness, fatigue, decreased exercise tolerance-- low ACTH, GH, LH, FSH, TSH
 - Weight loss--ACTH, DI
 - Increased body fat--GH, LH, FSH, TSH
 - Decreased muscle mass--GH, LH, FSH
 - Low libido, erectile dysfunction--LH, FSH
 - Ischemic heart disease--GH
 - Shortened life span--GH

Each deficiency gives multiple symptoms

	GH	TSH	ACTH	Gn	DI
• Poor growth	X	X			X
• Fatigue	X	X	X	X	
• Weight change	X	X	X		
• Drinking increase					X
• Precocious Puberty (child)				X	
• Loss of libido (adult)				X	

Review of post-TBI endocrine studies

- Types of published studies
 - Cases
 - Cross-sectional
 - All cases regardless of time since TBI
 - Prospective
 - Cases evaluated at baseline, then at intervals after TBI (such as 0, 3, 6, 12 mo after TBI)

Outline

- Mechanisms of injury
- Endocrine deficiencies
- Endocrine studies after TBI
 - Adult
 - Pediatric
- CCHMC study
- Consensus conference
- Conclusions

Cross-sectional studies in adults

	N	GH	ACTH	TSH	hi PRL	DI	LH, FSH
• 1988 Clark	53	100%	92%	88%	70%	26%	95%
• 2000 Bevenga	367	23	58	90	45	31	100
• 2001 Lieberman	70	15	7	22			
• 2004 Agha	102	11	13	1	12		12
• 2004 Bondanelli	50	28		10	8		14
• 2005 Leal-Cerro	113	6	6	6		2	17
• 2006 Herrmann	76	3	8	3	3		17
• 2006 Kelly	44	18					
• 2007 Bushnik	64	66	60	19			14
• 2008 Tarnivardi	29	21	6				
• 2009 Pavlovik	61	33	4.9	4.9			4.9
• 2010 Berg	246	5	4	12			9

Cross-sectional studies in adults

Overall N= 1275	Range %	Median %
• GHD	3 – 100	25
• Hypogonad	9 - 100	20
• TSHD	3 – 90	15
• High PRL	3 - 70	15
• DI	2 - 31	10
• ACTH-D	4 – 92	10

Meta-analysis

- Differences in incidence of deficiencies related to
 - Patient selection
 - All ICU admits vs referred, vs all with GHD
 - BMI
 - High BMI lowers GH peak
 - Definitions of deficiency
 - Testing methodology
 - Duration since TBI
 - Transient deficiencies in 1st y, chronic by 1y
- Kokshoorn 2010

Outline

- Mechanisms of injury
- Endocrine deficiencies
- Endocrine studies after TBI
 - Adult - Prospective
 - Pediatric
- CCHMC study
- Consensus conference
- Conclusions

Acute state after TBI

- 12 days after injury
 - 18% low GH to glucagon
 - 16% low cortisol to glucagon
 - 80% low gonadotropins
 - 52% high prolactin
 - 26% diabetes insipidus
 - 14% SIADH
 - 2% TSH deficiency


Agha 2004

Predictors of hypopituitarism

- 78 adults studied 3 & 12m after TBI
 - Diffuse axonal injury
 - Basal skull fracture
 - Older age

Schneider 2007


Change in pituitary function at 12 vs 3m after TBI



Aimaretti 2004

Prospective study, adult


Tanriverdi 2006



Outline

- Mechanisms of injury
- Endocrine deficiencies
- Endocrine studies after TBI
 - Adult
 - Pediatric
- CCHMC study
- Consensus conference
- Conclusions

Models of injury & recovery



Giza 2006

Case studies—Children after TBI

- 400 children (m/yr after head injury)
 - poor growth
 - Precocious puberty
 - Diabetes insipidus
 - failure to progress through puberty
- not known whether young brain/hypothalamus is more or less sensitive to injury than the older brain

Cross-sectional studies in children

	N	GH	ACTH	TSH	DI	LH,FSH
• 2006 Acerini (KIGS data)	141	100%	17%	17%		36%
• 2006 Einaudi	22	5	5			5
• 2007 Niederland	26	42	34	12		
• 2008 Poomthavorn	54	7	11	6	4	2

Cross-sectional studies in children

Overall N= 243	Range	Median
	%	%
• GHD	5 – 100	25
• Hypogonad	36	?
• Prec pub	2 - 5	?
• TSHD	6 - 17	12
• High PRL	-	?
• DI	4	?
• ACTH-D	5 – 34	20

Prospective study in older adolescents/young adults

- 3 & 12 mo after TBI
- 9 girls, 14 boys, age 16-25y
- At 3m, hypopituitarism in 35%
 - Total 9%, multiple 4%, isolated deficits 22%
 - Diabetes insipidus (DI) in 9%
 - mild hyperprolactinemia in 4%.
- At 12m, hypopituitarism in 30%
 - Total 9%, multiple 4%, isolated deficits 17%
 - DI in 4%
 - mild hyperprolactinemia in 4%

Aimaretti 2005

Prospective trial--children

- 30 acute, 26 at 6m, 20 at 12m
- At 12m
 - 3 with cerebral salt wasting
 - 1 DI
 - 1 with low cortisol
 - 1 with GHD

Einaudi 2006

Growth monitoring

- Prospective serial growth data was only obtained in only 22 of 129 children after TBI (requiring ICU care)
- Specialist & general pediatric followup after TBI should include growth measurements

Moon 2010

Outline

- Mechanisms of injury
- Endocrine deficiencies
- Endocrine studies after TBI
- CCHMC studies
 - Accidental TBI
 - Non-accidental TBI
- Consensus conference
- Conclusions

CCHMC prospective trial in children (support by Pfizer & GCRC)

- Authors & current location:
 - AMD Kaufers (Alabama), PF Backeljauw, K Reifschneider (Virginia), S Blum, L Michaud, M Weiss (New York), SR Rose
 - Several prior fellows made major contributions
 - Submitted for publication
- Objectives were to determine in children
 - incidence of endocrine abnormalities after TBI timing of endocrinopathies after TBI
 - best screen for central endocrinopathies after TBI

Methods

- Prospective observation
- N=31, mean age 11.6y (2-18y), 13 girls
- GCS ≤ 12
 - 12 had skull fracture (7 basilar)
 - 24 had intracranial hemorrhage
- Causes of injury
 - 15 motor vehicle crash
 - 7 fell
 - 7 hit by a car (4 riding a bike, 3 pedestrians)
 - 2 football injury
- recruited prior to hospital discharge after TBI
 - Consent, baseline medical history, physical exam, & laboratory testing

Glasgow Coma Scale

- 15 = normal
- 13 or 14+ mild head injury
- 9 - <13 moderate
- 3 - 8 severe
- ≤ 3 vegetative

Protocol Flow sheet

• Test/Procedure	At Injury	2-3	6-8m	12m
• Consent	X	X		
• History & Physical	X	X	X	X
• Serum/urine Osm.	X	X	X	
• AM Cortisol	X	X	X	X
• Free T4 and TSH	X	X	X	X
• IGF-I, IGFBP3	X	X	X	X
• Prolactin		X	X	repeat
• TSH surge			X	repeat
• 6 h GH secretion			X	stim tests
• 1 mcg ACTH test			X	repeat

Endocrinopathies after TBI

Endocrine abnl	Baseline (%)	2-3m (%)	6m (%)	12m (%)
Low thyroid	25	28	54	10
DI	7	4	--	--
Low GH	--	--	12	5
Puberty	--	20	17	19
Prolactin	--	12	33	--
Low ACTH	--	--	4	--
Any abnl	32	56	67	20

Figure 1





Figure 2



TBI severity did not predict endocrinopathy

	GCS	Skull Fracture	Intracranial Hemorrhage
Endocrinopathy at 12mo	6.7	43%	100%
Normal at 12mo	4.9	50%	92%

Conclusion from our study

- Similar to adults, many children have endocrine abnormalities persisting >6m after TBI
- Some abnormalities resolve by 1y after TBI
- Children <10yo may develop precocious puberty after TBI

Outline

- Mechanisms of injury
- Endocrine deficiencies
- Endocrine studies after TBI
- CCRM studies
 - Accidental TBI
 - Non-accidental TBI
- Consensus conference
- Conclusions

Children after inflicted injury in infancy

- No prior publications
- Cross sectional study of children with inflicted TBI
- IRB approved, informed consent obtained
- Inclusion criteria
 - inflicted TBI requiring hospitalization
 - injury at <3y of age
 - ≥ 1 y since injury
 - currently ≥ 2 to 9y of age
- 14 patients studied to date, ongoing

Methods

- Evaluation in overnight hospitalization
 - Height & weight
 - Height by stadiometer or length by infantometer
 - Morning cortisol & peak cortisol to low-dose cosyntropin
 - TSH nadir to peak rise at night (TSH surge) & free T4
 - 6h, every 20min GH sampling 10pm to 4am
 - hourly if <15kg
 - IGF-I, IGFBP3
 - Prolactin
 - Fasting serum & urine osmolality
 - LH, FSH, & estradiol or testosterone
 - If signs of puberty
- Patients grouped by prolactin level


Clinical Characteristics (according to Prolactin level)

Prolactin Group Mean \pm SD (ng/mL)	Gender	Average Age at Injury Mean \pm SD (months)	Time Elapsed at Study Mean \pm SD (months)	# Endocrine Abnormalities Mean \pm SD
High (n = 8) 25.7 \pm 12.3	6 M 2 F	10.6 \pm 11.5	36 \pm 21.6	2.6 \pm 0.7
Normal (n = 6) 10.8 \pm 6.7	5 M 1 F	3.2 \pm 3.4	36 \pm 8.4	0.8 \pm 0.8


Injury characteristics

- all patients had subdural hematoma
- most required ICU care
- 12 of 14 had ischemia or edema on CT


Acute injuries after iTBI (n = 14)




Height vs Age, High vs Normal Prolactin Groups



Pituitary Abnormalities according to Prolactin Group



Number of Pituitary Abnormalities



Results

- 86% (12 of 14)
 - at least 1 endocrine abnormality
- 64%
 - 2 or more abnormalities
- Elevated prolactin
 - most common, followed by short stature, abnormal thyroid function, & low GH peak
- High prolactin levels
 - associated with increased frequency of abnormal pituitary function
- Patients who required endocrine treatment are being followed clinically

Summary

- After inflicted TBI, many children have abnormal pituitary function
- If prolactin is elevated
 - need more complete hypothalamic-pituitary evaluation

Outline

- Mechanisms of injury
- Endocrine deficiencies
- Endocrine studies after TBI
- CCHMC study
- Consensus conference
- Conclusions

Consensus conference

- Better coordination needed among specialists
 - trauma surgeons, neurosurgeons, rehabilitation MDs, internists & endocrinologists
- Need to be aware of
 - risks for TBI-induced endocrinopathies
 - need for endocrine testing/ therapies

Ghigo 2005

Consensus conference

- Screening (8am lab) should include
 - cortisol
 - FT4, TSH
 - IGF-I
 - LH, FSH, T or E2
 - Prolactin
 - I's/ O's, serum sodium, urine & serum osmolality

Ghigo 2005

Recommendations

- After TBI in children
 - monitor length or height at least every 6 months
- Full pituitary function
 - evaluate at one year after injury
- Patients with elevated prolactin are likely to have other deficiencies

Conclusions

- Hypopituitarism may impair recovery after TBI
- Identification & therapy of deficiencies can improve chances of rehabilitation & enhance QoL
- Assessment of GH, ACTH, & TSH requires serial testing (re-eval at 1y)
- Need close liaison between endocrine, rehabilitation, & neurosurgical services

